CALYCANTHUS FLORIDUS (CALYCANTHACEAE)— A NOMENCLATURAL NOTE

D. E. BOUFFORD AND S. A. SPONGBERG

In his revisionary study of the Calycanthaceae, Nicely (1965) recognized a single eastern North American species of Calycanthus, which comprises two varieties, C. floridus L. var. floridus and C. floridus var. laevigatus (Willd.) Torrey & Gray. This treatment has been either adopted independently or followed by Radford, Ahles, and Bell (1964, 1965, 1968) and Strausbaugh and Core (1978), and subsequently by most, if not all, collectors in the southeastern United States. Although Nicely's treatment may oversimplify the variation patterns present in Calycanthus in the eastern United States (cf. Wood, 1958), we are not prepared at this time to refute his taxonomic scheme. However, we should like to draw attention to the fact that laevigatus is not the earliest varietal epithet available when Nicely's treatment of C. floridus is followed.

Torrey and Gray (1840) were the first to use *laevigatus* as a varietal epithet under *Calycanthus floridus* when they reduced *C. laevigatus* Willd., *C. glaucus* Willd., and *C. inodorus* Ell. to varietal standing. Since Nicely's circumscription of plants of *C. floridus* with glabrous lower leaf surfaces includes both glaucous and nonglaucous forms, it appears that his choice of *laevigatus* (rather than *glaucus* or *inodorus*) as the varietal epithet was based on a desire to use an all-inclusive, descriptive epithet.

Nuttall (1818), however, recognized four taxa of Calycanthus in eastern North America—three species (C. floridus, C. glaucus, and C. laevigatus) and one variety (C. glaucus var. oblongifolius Nutt.). Since Nicely considered C. glaucus var. glaucus, C. glaucus var. oblongifolius, C. floridus var. glaucus, and C. floridus var. laevigatus (as well as C. floridus var. inodorus) to be synonyms that apply to the same taxon, Nuttall's epithet oblongifolius must be considered when determining the correct name of the variety of C. floridus. Oblongifolius (1818) predates Torrey and Gray's usage of glaucus, inodorus, and laevigatus (all 1840) as varietal epithets by 22 years and therefore has priority as the earliest available varietal epithet for plants of C. floridus with glabrous leaf undersurfaces as circumscribed by Nicely. For this reason the following new combination is necessitated.

Calycanthus floridus L. var. oblongifolius (Nutt.) D. E. Boufford & S. A. Spongberg, comb. nov.

Article 60 of the International Code of Botanical Nomenclature (Regnum Vegetabile 97: 51. 1978) states that "when the rank of a genus or infrageneric taxon is changed, the correct name or epithet is the earliest legitimate one available in the new rank. In no case does a name or epithet have priority outside its own rank."

[©] President and Fellows of Harvard College, 1981.

Journal of the Arnold Arboretum 62: 265, 266. April, 1981.

BASIONYM: Calycanthus glaucus Willd. var. oblongifolius Nutt. Gen. N. Am. Pl. 1: 312. 1818.

Synonyms, which should include Calycanthus floridus var. laevigatus, can be found in Nicely (1965).

REFERENCES

- NICELY, K. A. A monographic study of the Calycanthaceae. Castanea 30: 38-81. 1965.
- NUTTALL, T. The genera of North American plants 1: 311, 312. 1818.
- RADFORD, A. E., H. E. Ahles, & C. R. Bell. Guide to the vascular flora of the Carolinas. 383 pp. The Book Exchange, Univ. North Carolina, Chapel Hill. 1964.
- pp. North Carolina Agr. Exper. Sta. Tech. Bull. No. 165, 1965.
- Strausbaugh, P. D., & E. L. Core. Flora of West Virginia. ed. 2. x1 + 1079 pp. Seneca Books, Inc., Grantsville, West Virginia. 1978.
- Wood, C. E., Jr. The genera of the woody Ranales in the southeastern United States. Jour. Arnold Arb. 39: 296-346. 1958.
 - D. E. B.
 Carnegie Museum of Natural
 History
 4400 Forbes Avenue
 Pittsburgh, Pennsylvania 15213
- S. A. S.
 THE ARNOLD ARBORETUM OF
 HARVARD UNIVERSITY
 22 DIVINITY AVENUE
 CAMBRIDGE, MASSACHUSETTS
 02138